

Density expectation value of two independent interacting Bose-Einstein condensates

Hongwei Xiong^{1,2,3}, Shujuan Liu^{1,2,3}, Mingsheng Zhan^{1,2}

¹State Key Laboratory of Magnetic Resonance and Atomic and Molecular Physics,
Wuhan Institute of Physics and Mathematics, Chinese Academy of Sciences, Wuhan 430071, P. R. China

²Center for Cold Atom Physics, Chinese Academy of Sciences, Wuhan 430071, P. R. China and

³Graduate School of the Chinese Academy of Sciences, P. R. China

(Dated: February 2, 2008)

After removing the double-well potential trapping two initially independent Bose condensates, the density expectation value is calculated when both the exchange symmetry of identical bosons and interatomic interaction are considered. After the overlapping, it is shown that there is a nonzero interference term in the density expectation value. This nonzero interference term physically arises from the exchange symmetry of identical bosons and interatomic interaction which make two initially independent condensates become coherent after the overlapping. It is found that the calculated density expectation value with this model agrees with the interference pattern observed in the experiment by Andrews *et al* (*Science* **275**, 637 (1997)).

PACS: 05.30.Jp; 03.75.Kk; 03.65.Ta

The coherence property plays an essential role in the wave nature of Bose-Einstein condensates (BECs), and it has been investigated intensively [1] after the experimental realizations of BECs in dilute gases. For two coherently separated Bose condensates, it is not surprising that there is a clear interference pattern when the two condensates are allowed to overlap. In the celebrated experiment by Andrews *et al* [2], however, high-contrast fringes were observed even for two completely independent condensates at an initial time. This experimental result shows clearly that there is a coherence property after the overlapping of two independent interacting condensates.

In many literature (see for example [3, 4] and references therein), upon expansion, it is shown that there is no interference term in the density expectation value for two initially independent condensates, and thus the observed high-contrast fringes [2] for two independent condensates were interpreted [3] with the aid of the high-order correlation function $P(\mathbf{r}, \mathbf{r}', t)$ (which is an oscillation function of $\mathbf{r} - \mathbf{r}'$) and quantum measurement process. The contradiction between the vanishing interference term in the density expectation value and the observed high-contrast fringes was also discussed based on several theories such as the stochastic simulations of the photon detection for atoms [5], the expansion of Fock state by the linear superposition of coherent states [6], and the continuous measurement theory [7]. In addition, the nonlinear effects in the interference pattern of two coherently separated Bose condensates were investigated in Refs [8, 9].

In the present work, we calculate carefully the density expectation value for two initially independent condensates by including the interatomic interaction. It is found that *for the case of two initially independent condensates, upon expansion, there is a nonzero interference term in the density expectation value when interatomic interaction and the exchange symmetry of identical bosons are taken into account carefully!* After removing the double-well potential trapping the two initially independent condensates, we give the theoretical result of the density ex-

pectation value which agrees with the interference pattern observed in [2].

For the condensate trapped in a magnetic trap, it can be divided into two condensates by a blue-detuned laser beam. The combined double-well potential takes the form

$$V_{ext} = \frac{1}{2}m \left[\omega_x^2 (x - x_0)^2 + \omega_y^2 y^2 + \omega_z^2 z^2 \right] + U_0 e^{-(x-x_0)^2/w_x^2 - (y^2+z^2)/w_\perp^2}, \quad (1)$$

where the second term represents the external potential due to the laser beam.

After the evaporative cooling for Bose gases trapped in the double-well potential, if the intensity of the blue-detuned laser beam is sufficiently low so that the tunneling effect is obvious, the two condensates can be regarded to be coherently separated. For two coherently separated condensates, every atom is described by the following normalization wavefunction

$$\phi_c(\mathbf{r}, t) = \left[\sqrt{N_1} \phi_1(\mathbf{r}, t) + \sqrt{N_2} \phi_2(\mathbf{r}, t) \right] / \sqrt{N}, \quad (2)$$

where $\phi_1(\mathbf{r}, t)$ and $\phi_2(\mathbf{r}, t)$ are the normalization wave functions accounting for the two condensates. $N = N_1 + N_2$ is the total number of particles. Before the overlapping of the two condensates, the average particle numbers in each condensate are N_1 and N_2 , respectively. After removing the double-well potential, the density expectation value is then

$$n_c(\mathbf{r}, t) = N \left[a_c |\phi_1(\mathbf{r}, t)|^2 + 2b_c \times \text{Re}(\phi_1^*(\mathbf{r}, t) \phi_2(\mathbf{r}, t)) + c_c |\phi_2(\mathbf{r}, t)|^2 \right], \quad (3)$$

where $a_c = N_1/N$, $b_c = \sqrt{N_1 N_2}/N$ and $c_c = N_2/N$. The second term in the above equation accounts for the interference effect when there is an overlapping between two condensates upon expansion.

If the intensity of the blue-detuned laser beam is sufficiently high so that the tunneling effect can be omitted,

the two condensates can be regarded to be completely independent. In this case, the number of particles N_1 and N_2 in each of the two condensates is fixed [4]. To investigate clearly the role of the exchange symmetry of identical particles and interatomic interaction, we calculate the density expectation value $n_d(\mathbf{r}, t)$ directly from the many-body wave function. Based on the well-known exchange symmetry of identical bosons [10], the many-body wavefunction of the whole system is given by

$$\begin{aligned} & \Psi_{N_1 N_2}(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N, t) \\ &= A_n \sqrt{\frac{N_1! N_2!}{N!}} \sum_P P [\phi_1(\mathbf{r}_1, t) \cdots \phi_1(\mathbf{r}_{N_1}, t) \times \\ & \quad \phi_2(\mathbf{r}_{N_1+1}, t) \cdots \phi_2(\mathbf{r}_{N_1+N_2}, t)], \end{aligned} \quad (4)$$

where P denotes the $N!/(N_1! N_2!)$ permutations for the particles in different single-particle state ϕ_1 or ϕ_2 . The above expression of the many-body wavefunction is based on the quantum mechanical principle [10]. A_n is a normalization factor to assure $\int |\Psi_{N_1 N_2}(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N, t)|^2 d\mathbf{r}_1 d\mathbf{r}_2 \cdots d\mathbf{r}_N = 1$. A_n is determined by the following equation:

$$A_n \left[\sum_{i=0}^{\min(N_1, N_2)} \frac{N_1! N_2! |\zeta(t)|^{2i}}{i! i! (N_1 - i)! (N_2 - i)!} \right]^{1/2} = 1, \quad (5)$$

where

$$\zeta(t) = \int \phi_1(\mathbf{r}, t) \phi_2^*(\mathbf{r}, t) dV = |\zeta(t)| e^{i\varphi_c}. \quad (6)$$

In this Letter, to give a concise expression for various coefficients such as A_n , we have introduced the rule $0^0 = 1$.

From the above many-body wavefunction, the exact density expectation value takes the following form:

$$\begin{aligned} n_d(\mathbf{r}, t) &= N \int \Psi_{N_1 N_2}^*(\mathbf{r}, \mathbf{r}_2, \dots, \mathbf{r}_N, t) \times \\ & \quad \Psi_{N_1 N_2}(\mathbf{r}, \mathbf{r}_2, \dots, \mathbf{r}_N, t) d\mathbf{r}_2 \cdots d\mathbf{r}_N \\ &= N B_n \left[a_d |\phi_1|^2 + 2b_d \times \text{Re}(e^{i\varphi_c} \phi_1^* \phi_2) \right. \\ & \quad \left. + c_d |\phi_2|^2 \right], \end{aligned} \quad (7)$$

where $B_n = A_n^2 N_1! N_2! / N!$, and the coefficients in the above equation are given by

$$a_d = \sum_{i=0}^{\min(N_1-1, N_2)} \frac{(N-1)! |\zeta(t)|^{2i}}{i! i! (N_1 - i - 1)! (N_2 - i)!}, \quad (8)$$

$$\begin{aligned} b_d &= \sum_{i=0}^{\min(N_1-1, N_2-1)} \frac{(N-1)! |\zeta(t)|^{2i+1}}{i! (i+1)! (N_1 - i - 1)! (N_2 - i - 1)!}, \quad (9) \\ c_d &= \sum_{i=0}^{\min(N_1, N_2-1)} \frac{(N-1)! |\zeta(t)|^{2i}}{i! i! (N_1 - i)! (N_2 - i - 1)!}. \quad (10) \end{aligned}$$

For two independent ideal condensates, before the overlapping of the two condensates, $\zeta(t=0) = 0$. Based on the Schrödinger equation, it is easy to verify that after the double-well potential separating the condensates is removed, one has $\zeta(t) = 0$ at any further time. Thus $b_d = 0$, and the density expectation value is given by

$$n_d(\mathbf{r}, t) = N_1 |\phi_1(\mathbf{r}, t)|^2 + N_2 |\phi_2(\mathbf{r}, t)|^2. \quad (11)$$

In this situation, the interference term is zero in the density expectation value.

However, if the interatomic interaction is considered, $\zeta(t)$ can be a nonzero value, and this will lead to a quite different result compared with the ideal condensates. Assuming that $g = 4\pi\hbar^2 a_s / m$ with a_s being the scattering length, after removing the double-well potential, the energy of the whole system is given by

$$\begin{aligned} E &= \int \Psi_{N_1 N_2}^* \left[\sum_{i=1}^N -\frac{\hbar^2}{2m} \nabla_i^2 \right] \Psi_{N_1 N_2} d\mathbf{r}_1 \cdots d\mathbf{r}_N \\ &+ g \int \Psi_{N_1 N_2}^* \sum_{i < j}^N \delta(\mathbf{r}_i - \mathbf{r}_j) \Psi_{N_1 N_2} d\mathbf{r}_1 \cdots d\mathbf{r}_N. \end{aligned} \quad (12)$$

The second term in the above equation represents the interaction energy of the whole system by using the well-known two-body pseudopotentials.

By using the ordinary action principle and the energy of the whole system, one can get the following coupled evolution equations for ϕ_1 and ϕ_2 :

$$i\hbar \frac{\partial \phi_1}{\partial t} = \frac{1}{N_1} \frac{\delta E}{\delta \phi_1^*}, \quad (13)$$

$$i\hbar \frac{\partial \phi_2}{\partial t} = \frac{1}{N_2} \frac{\delta E}{\delta \phi_2^*}, \quad (14)$$

where $\delta E / \delta \phi_1^*$ and $\delta E / \delta \phi_2^*$ are functional derivatives. With the above coupled equations, one can understand that $\zeta(t)$ becomes nonzero after the overlapping between two initially independent condensates for g being a nonzero value [11].

Although generally speaking, $|\zeta(t)|$ is much smaller than 1 because $\phi_1 \phi_2^*$ is an oscillation function about the space coordinate, nevertheless, a nonzero value of $\zeta(t)$ will give significant contribution to the density expectation value for large N_1 and N_2 . In Fig. 1(a), we give the relation between b_d/c_d and ζ for $N_1 = N_2 = 10^3$ based on the numerical calculations of Eqs. (8), (9) and (10). The relation between b_d/c_d and $N_1 = N_2$ for $\zeta = 0.001$ is shown in Fig. 1(b). It is shown clearly that even for $|\zeta|$ being the order of N_1^{-1} , the value of b_d can not be omitted, and thus the interference term in Eq. (7) will play an important role in the density expectation value. Generally speaking, for $N_1 |\zeta| \gg 1$ and $N_2 |\zeta| \gg 1$, one has $b_d / \sqrt{a_d c_d} \approx 1$.

Generally speaking, increasing the particle number will enhance the effect of the interference term in the density




FIG. 1: Based on the numerical calculations of Eqs. (8), (9) and (10), shown in Fig. 1(a) is the relation between b_d/c_d and ζ , while shown in Fig. 1(b) is the relation between b_d/c_d and $N_1 = N_2$ for $\zeta = 0.001$. It is shown clearly that the interference term can play an important role in the density expectation value for $|\zeta|$ being larger than N_1^{-1} .

expectation value. Based on Eqs. (13) and (14), increasing the coupling constant g will have the effect of increasing $\zeta(t)$. Together with the relation between b_d/c_d and ζ illustrated in Fig. 1(a), this shows that increasing the interatomic interaction will increase the effect of the interference term.

In the experiment of Ref. [2], $N = 5 \times 10^6$ condensed sodium atoms were confined in a magnetic trap with $\omega_x = 2\pi \times 18$ Hz, $\omega_y = \omega_z = 2\pi \times 320$ Hz. A blue-detuned laser beam of wavelength 514 nm was focused into a light sheet with a cross section of $12 \mu\text{m}$ by $67 \mu\text{m}$. The long axis of the laser beam was perpendicular to the long x -axis of the condensate. For a laser power of 14 mW, the barrier height is about $1.4 \mu\text{K}$, which is much larger than the chemical potential $\mu = 0.03 \mu\text{K}$. For this laser beam, the two condensates can be regarded to be independent because they are well separated and the tunneling effect can be omitted (See also [3]). With these experimental parameters and the s -wave scattering length $a_s = 2.75$ nm, the initial profile of the two condensates is shown in Fig. 2. After the double-well potential is removed, the evolution of the density expectation value $n_{d-x}(x, t) = \int n_d(\mathbf{r}, t) dy dz$ (in unit of $N/2$) is given in Fig. 2 through the numerical calculations of Eqs. (7), (13) and (14). We see that there is a clear interference pattern in the density expectation value which agrees with the experimental result. Shown in the inset is the evolution of $|\zeta|$ for these parameters. We have also verified in the numerical calculation that $|\zeta|$ is always zero for two ideal condensates. For the expansion time of 40 ms, the numerical result of $|\zeta|$ shows that $b_d/c_d \approx 1$.

Based on the evolution of the density expectation value shown in Fig. 2, we see that the overall width (about




FIG. 2: Shown is the evolution of the density expectation value calculated using the experimental parameters in Ref. [2]. It is shown that there is a clear interference pattern after the overlapping of two independent interacting condensates. Shown in the inset is the evolution of $|\zeta(t)|$ for the experimental parameters in Ref. [2].

$300 \mu\text{m}$) of the ultracold gases in x -direction does not increase obviously. This is due to the fact that the initial density distribution is cigar-shaped, and thus the expansion in x -direction is very slow, while the expansion in y and z directions is much quick. In the experiment of Ref. [12], one can see clearly that there is no obvious expansion in the long x -axis for cigar-shaped condensate. For two initially independent condensates, when the double-well potential is switched off, one should note that in the regime close to $x = 250 \mu\text{m}$ shown in Fig. 2, the ultracold gases expand rapidly in x -direction because in this regime the ultracold gases have higher kinetic energy. Thus, although the total width of the system in x -direction does not increase obviously, the rapid expansion in the central regime will lead to the overlapping between two initially independent condensates, and results in the interference effect. The overall width of the system shown in Fig. 2 is smaller than the experimental result of about $500 \mu\text{m}$ [2]. This difference may come from the expansion of thermal cloud in this experiment [2]. After 40 ms expansion, the numerical result in [8] for two coherently separated condensates also showed that the overall width of the system is about $300 \mu\text{m}$.

The essential reason for the emergence of the interference term of two initially independent condensates lies in that because of the exchange symmetry of identical bosons and interatomic interaction, the two initially independent condensates will become coherent after the overlapping between the two condensates. The physical mechanism of this interaction-induced coherence can be also understood based on the second quantization method. For two initially independent condensates, the

quantum state of the whole system can be written as [4]

$$|N_1, N_2\rangle = \frac{C_n}{\sqrt{N_1!N_2!}} (\hat{a}_1^\dagger)^{N_1} (\hat{a}_2^\dagger)^{N_2} |0\rangle, \quad (15)$$

where C_n is a normalization constant to assure $\langle N_1, N_2 | N_1, N_2 \rangle = 1$. $\hat{a}_1^\dagger$ ($\hat{a}_2^\dagger$) is a creation operator which creates a particle described by the single-particle state ϕ_1 (ϕ_2) in the left (right) condensate. The density expectation value is then

$$n_d(\mathbf{r}, t) = \langle N_1, N_2, t | \hat{\Psi}^\dagger \hat{\Psi} | N_1, N_2, t \rangle. \quad (16)$$

Here $\hat{\Psi}(\mathbf{x}, t)$ is the field operator. The operators $\hat{a}_1$ and $\hat{a}_2$ can be written as $\hat{a}_1 = \int \hat{\Psi} \phi_1^* dV$ and $\hat{a}_2 = \int \hat{\Psi} \phi_2^* dV$, respectively. By using the commutation relations of the field operators $[\hat{\Psi}(\mathbf{x}, t), \hat{\Psi}(\mathbf{y}, t)] = 0$ and $[\hat{\Psi}(\mathbf{x}, t), \hat{\Psi}^\dagger(\mathbf{y}, t)] = \delta(\mathbf{x} - \mathbf{y})$. It is easy to get the commutation relation $[\hat{a}_1, \hat{a}_2^\dagger] = \zeta^*$. We see that $\hat{a}_1$ and $\hat{a}_2^\dagger$ are not commutative any more for $\int \phi_1 \phi_2^* dV$ being a nonzero value. The non-commutative property between $\hat{a}_1$ and $\hat{a}_2^\dagger$ plays an essential role in the emergence of the interference term. Our calculations show that the result given by Eq. (16) is the same as the result given by Eq. (7) which is obtained based on the many-body wavefunction. We have also proven that the evolution equations for ϕ_1 and ϕ_2 based on the many-body wavefunction are the same as the results based on the second quantization method [11].

One can get a further physical picture for the emergence of interference pattern of two initially independent condensates through a general investigation for the case of $N_1 |\zeta| \gg 1$, $N_2 |\zeta| \gg 1$ and $N_1 \sim N_2$. Based on our numerical calculations, these conditions are satisfied for the case of 40 ms expansion in the experiment of Ref. [2]. When these conditions are satisfied, $NB_n a_d \approx N_1$, $NB_n b_d \approx \sqrt{N_1 N_2}$ and $NB_n c_d \approx N_2$. Thus, the density expectation value can be approximated very well as

$$n_d(\mathbf{r}, t) \simeq \Phi_e^*(\mathbf{r}, t) \Phi_e(\mathbf{r}, t), \quad (17)$$

where the effective order parameter $\Phi_e(\mathbf{r}, t)$ is

$$\Phi_e(\mathbf{r}, t) = \sqrt{N_1} \phi_1(\mathbf{r}, t) + \sqrt{N_2} e^{i\varphi_c} \phi_2(\mathbf{r}, t). \quad (18)$$

In particular, after removing the double-well potential, the evolution of the effective order parameter $\Phi_e(\mathbf{r}, t)$ can be described very well by the following Gross-Pitaevskii equation [11]

$$i\hbar \frac{\partial \Phi_e}{\partial t} \simeq -\frac{\hbar^2}{2m} \nabla^2 \Phi_e + g |\Phi_e|^2 \Phi_e. \quad (19)$$

We see that the emergence of the effective order parameter gives us strong evidence that the coherence is formed between two initially independent condensates, and thus results in the emergence of high-contrast interference fringes.

In summary, upon expansion, we calculate the density expectation value of two initially independent condensates. Quite different from the current beliefs, it is found that there is a nonzero interference term when the interatomic interaction and the exchange symmetry of identical bosons are both considered carefully. In fact, it is well-known that the interaction plays an essential role in the formation of the order parameter of Bose-condensed gases, i.e. the formation of a stable coherent property. Here, we provide an example in which the interaction induces coherent evolution between two initially independent condensates. To investigate more clearly the interference pattern due to the measurement process alone, an experimental investigation of two independent ideal condensates would be very interesting, because there is no interference term for ideal condensates in the density expectation value. In the last few years, the rapid experimental advances of Feshbach resonance where the scattering length can be tuned from positive to negative make this sort of experiment be feasible.

-
- [1] P. Meystre, *Atom optics* (Springer-Verlag, New York, 2001).
 - [2] M. R. Andrews, C. G. Townsend, H. -J. Miesner, D. S. Durfee, D. M. Kurn, and W. Ketterle, *Science* **275**, 637 (1997).
 - [3] A. J. Leggett, *Rev. Mod. Phys.* **73**, 307 (2001).
 - [4] C. J. Pethick and H. Smith, *Bose-Einstein Condensation in Dilute Gases* (Cambridge University, Cambridge, 2002).
 - [5] J. Javanainen and S. M. Yoo, *Phys. Rev. Lett.* **76**, 161 (1996).
 - [6] Y. Castin and J. Dalibard, *Phys. Rev. A* **55**, 4330 (1997).
 - [7] J. I. Cirac, C. W. Gardiner, M. Naraschewski, P. Zoller, *Phys. Rev. A* **54**, R3714 (1996).
 - [8] A. Röhrl, M. Naraschewski, A. Schenzle, and H. Wallis, *Phys. Rev. Lett.* **78**, 4143 (1997).
 - [9] W. M. Liu, B. Wu, and Q. Niu, *Phys. Rev. Lett.* **84**, 2294 (2000).
 - [10] L. D. Landau and E. M. Lifshitz, *Quantum mechanics* (Pergamon, Oxford, 1977).
 - [11] Detailed analyses will be given elsewhere.
 - [12] M.-O. Mewes, M. R. Andrews, N.J. van Druten, D. M. Kurn, D. S. Durfee, and W. Ketterle, *Phys. Rev. Lett.* **77**, 416 (1996).